

Three novel constituents from the roots of *Rhaponticum carthamoides*

ZOLTÁN PÉTER ZOMBORSZKI¹, NORBERT KÚSZ¹, JUDIT HOHMANN^{1,2}, DEZSŐ CSUPOR^{1,3,*}

¹Institute of Pharmacognosy, Faculty of Pharmacy, University of Szeged, H-6720 Szeged, Eötvös u. 6, Hungary

²Institute of Clinical Pharmacy, Faculty of Pharmacy, University of Szeged, H-6725 Szeged, Szikra u. 8, Hungary

³ELKH-USZ Biologically Active Natural Products Research Group, University of Szeged, H-6720 Szeged, Eötvös u. 6, Hungary

*Corresponding author: Dezső Csupor

Email: csupor.dezso@szte.hu

Received: 15 January 2023 / Revised: 2 February 2023 / Accepted: 3 February 2023

Aims: The purpose of our study was to conduct a phytochemical analysis of *Rhaponticum carthamoides* in order to gain a better understanding of its chemical profile and the active constituents responsible for its bioactivities.

Methods: The methanolic extract of the root was fractionated using multiple chromatographic methods, including normal, medium and high-pressure chromatography, rotation planar chromatography with a variety of stationary phases and eluents. The structures of the isolated compounds were identified by comparing their NMR spectra with those reported in the literature.

Results: As a result of our experiments, the mixture of *cis* and *trans* isomers of tiophene derivatives, vanillic acid, chlorogenic acid methyl ester, ajugasterone C, makisterone C, tachioside, 20-hydroxyecdysone, 24-epi-makisterone A and evofolin B were isolated and identified in the root of the plant.

Conclusion: Tachioside, evofolin B and chlorogenic acid methyl ester have been reported for the first time from this species.

Keywords: *Rhaponticum carthamoides*, *Leuzea carthamoides*, tachioside, evofolin B, chlorogenic acid methyl ester

1 Introduction

Rhaponticum carthamoides (Willd) Iljin, also known as *Leuzea carthamoides* (Willd) DC. or Maral root is a perennial plant belonging to the Asteraceae family. It is indigenous in the subalpine zone of Altay and Sayan Mountains, Siberia, Kazakhstan and Eastern Europe. The plant has been used in the folk medicine for its beneficial properties and thus widely cultivated, especially in Russia. Its main therapeutic indication is based on the adaptogenic effects of the roots and it is used as a general tonic and anabolic to enhance physical and mental performance in physical workers and athletes, and to promote recovery after illnesses and surgical interventions (1–6).

The main constituents of *R. carthamoides* belong to phytoecdysteroids, which are playing a defensive role against insects, acting as inhibitors of insect growth and moulting. In the last decades more than 50 ecdysteroids and their derivatives have been isolated from the plant, the majority of these compounds being identified in the roots, but the seeds and leaves also contain this type of substances (2,7–17). The most abundant constituent is 20-hydroxyecdysone, also known as β -ecdysone or polypodine A. The adaptogenic effect of the

plant can be explained by the presence of ecdysteroids. Besides phytoecdysteroids, flavonoids, phenolic acids, lignans, polyacetylenes, sesquiterpene lactones and triterpenes have also been detected from this species (18).

Because of the adaptogenic properties of the *R. carthamoides*, numerous studies were carried out with the plant, aiming to understand its pharmacology. Various animal studies have been performed with the extracts of the plant and its main ecdysteroid, 20-hydroxyecdysone. Increase in protein synthesis and body mass were observed in different experiments, moreover, increased work capacity was also reported (19–23). The adaptogenic effect was studied in human trials as well. In clinical studies with athletes, significant muscle mass increase was observed in the groups treated with *R. carthamoides* or 20-hydroxyecdysone compared with the control (19,24).

However, the pharmacological profile of *R. carthamoides* goes beyond its anabolic effects. The plant extract exerts immunomodulatory activities, which may be explained by the presence of other compounds than ecdysteroids (25,26). The same applies to its central nervous system (27–29) and hemorheological effects (30).

Considering the perspective use of *R. carthamoides* in the medicine and its undiscovered phytochemical profile, our aim was to identify further compounds from this plant.

2 Materials and methods

2.1 General experimental procedures

The HPLC system comprised of Waters 600 pump, Waters In-line degasser AF, Waters 2487 dual channel UV detector modules connected with Waters 600 control module. Rotation planar chromatography (RPC) was performed with a Chromatron instrument (model 8924; Harrison Research) on manually coated SiO₂ plates.

The isolated compounds were analysed via NMR. NMR spectra (Supplementary material) were recorded mostly in MeOH-*d*₄ for ¹H spectra, but DMSO-*d*₆ and CDCl₃ were also used for 2D on a Bruker Avance 600 III spectrometer (¹H: 600.13 MHz; ¹³C: 150.9 MHz) equipped with a 5 mm cryo-TXI probe. The peaks of the residual solvents were taken as reference points. Chemical shifts are expressed in parts per million and coupling constants (*J*) values are reported in Hz. Data were acquired and processed with the MestReNova

v6.0.2-5475 software and were compared to literature data to identify the compounds.

2.2 Plant material

Rhaponticum carthamoides roots ("Roots of *Leuzea carthamoides* [Lujza]" 2,0 kg) were purchased from Herbosus (Finland, Espoo). The plant material was identified by dr. Zsuzsanna Hajdú (University of Szeged, Institute of Pharmacognosy).

2.3 Extraction and isolation

The dry roots (1.75 kg) were ground, then extracted with MeOH, using an ultrasonic bath at room temperature. The extract then was filtered and was evaporated to dryness, yielding 112.52 grams of dry extract. To the dry residue 1000 mL of MeOH – H₂O (1:1) was added, then extracted with *n*-hexane, CH₂Cl₂ and EtOAc, resulting dry fractions 9.91 g, 28.79 g and 103.37 g, respectively (Figure 1). The EtOAc fraction was fractionated by column chromatography on SiO₂ stationary phase with a gradient elution using CH₂Cl₂ – MeOH (95:5–0:100). After TLC analysis the fractions were merged to subfractions A–T. Purification of subfraction E by rotation planar chroma-




Figure 1 Isolation of metabolites from the roots of *R. carthamoides*

tography (RPC) on 4 mm SiO₂ plates using an *n*-hexane – EtOAc gradient (100:0–0:100) led to sub-fractions **EI–EXXI**. With the help of HPLC fraction **EX** was further purified on a Phenomenex Kinetex C18 column (250×21.1 mm, 100 Å, 5 μm) using H₂O – MeOH (0 min: 40:60, 15 min: 0:100, 16 min: 0:100, 20 min: 40:60) as eluent. The resulting fractions were further refined with atmospheric pressure column using SiO₂ as stationary and CH₂Cl₂ – EtOAc [100:0–0:100, 10% (20ml each) increments] as mobile phase. As a result, the pure compound **1** (21 mg) were obtained. The subfraction **EXX** was further purified using RPC [1 mm SiO₂ CH₂Cl₂ – EtOH (100:0–0:100, 3.0 mL/min)] and preparative TLC (SiO₂, elution with EtOAc – EtOH – H₂O (4:0.5:0.25) to obtain the pure compound **2** (13 mg).

Subfraction **L** was fractionated with help of medium pressure liquid chromatography (MPLC) using SiO₂ (0.043–0.060 mm) as stationary phase eluted with CH₂Cl₂ – EtOAc gradient (100:0–0:100). After TLC analysis the gained fractions were combined to five subfractions, namely **LI–LV**. The **LII** was further processed on an atmospheric column filled with SiO₂ (0.043–0.060 mm). EtOAc – MeOH (100:0–0:100) was used as eluent. With a subsequent purification using HPLC [LiChrospher RP-C18e column (250×4.6mm, 5 μm), gradient elution with H₂O – MeOH (90:10–50:50) in fifteen minutes], one pure compound (**3**, 10 mg) was obtained. The subfraction **LIV** was further purified by MPLC. As stationary phase SiO₂ (0.043–0.060 mm) was chosen. The gradient elution consisted of EtOAc – MeOH (0 min: 100:0, 60 min: 90:10, 70




Figure 2 Compounds 1–9 Isolated from *R. carthamoides*

min: 90:10). A further fractionation by HPLC (Li-Chrospher RP-C18e, 250×4.6 mm, 5 µm column, H₂O – MeOH gradient) resulted in the isolation of the pure compounds **4** (8 mg), **5** (2 mg), **6** (2 mg), **7** (43 mg) and **8** (2 mg).

The fraction **F** was further processed by MPLC. To this SiO₂ (0.043–0.060 mm) as stationary phase was used. As eluent hexane – CH₂Cl₂ – MeOH (100:0:0–0:100:0–0:50:50), with a flow rate of 15 mL/min was utilised. After TLC analysis and combination of fractions the resulting subfractions were **FI–IX**. The fraction **FVIII** was further purified on MPLC on SiO₂ stationary phase (0.043–0.060 mm) using a gradient elution of CH₂Cl₂ – MeOH – H₂O (100:0:0–100:0:0 to 0:50:50), with a flow rate of 25 mL/min with 5% increments. A final purification with HPLC (Kinetex Phenomenex C18, 150×4.6 mm, 5 µm; H₂O – MeCN gradient) resulted in the isolation of the pure compound **9** (5 mg).

3 Results

As result of a series of various chromatographic purifications, such as normal, medium and high-pressure chromatography, rotation planar chromatography with a variety of stationary phases and eluents, combined with modern structure elucidation methods (NMR, spectra see in Supplementary material) in total nine compounds were isolated from the plant. These compounds were identified by comparing their NMR spectra with those reported in the literature as the mixture of *cis* and *trans* isomers of thiophene derivatives (compound **1**) (31) vanillic acid (**2**) (32), chlorogenic acid methyl ester (**3**) (33), ajugasterone C (**4**) (34), makisterone C (**5**) (35) tachioside (**6**) (36), 20-hydroxyecdysone (**7**) (37) 24-epi-makisterone A (**8**) (35), evofolin B (**9**) (38) (Figure 2).

4 Discussion

Thiophene derivatives (31), vanillic acid (39), ajugasterone C (40), makisterone C (41), 20-hydroxyecdysone (41), 24-epi-makisterone A (35) have been previously reported from this plant, however, the presence of tachioside, evofolin B and chlorogenic acid methyl ester have been reported from *R. carthamoides* for the first time by our research group.

The ecdysteroids ajugasterone C, makisterone C, 20-hydroxyecdysone and 24-epi-makisterone contribute to the adaptogenic effect of the plant. The contribution of other compounds than ecdysteroids in the clinical effects of the plant is slightly

discovered. The specific role of vanillic acid and chlorogenic acid methyl ester is not likely, since these compounds are ubiquitous in the plant kingdom and were isolated in small amounts from the plant. Evofolin B, a benzenoid isolated for the first time in 1995 (42), has received little pharmacological attention. The weak *in vitro* quinone reductase inducing (43), the *in vitro* superoxide anion generation inhibitory (44) the moderate *in vitro* lipolytic (45) activities are not directly related to the therapeutic use of *R. carthamoides*. Tachioside, an aromatic glycoside that was first isolated from *Berchemia racemosa* (46) is reported to have *in vitro* tyrosinase (47), moderate alpha-glucosidase (48) and 15-lipoxygenase inhibitory activities (49) and possesses antioxidant effect (50). However, some of the reported activities of these compounds make them interesting for future investigations and could also open up new opportunities for *R. carthamoides* research.

5 Conclusions

As the result our experiments we isolated and characterised three, previously undescribed compounds (tachioside, evofolin B and chlorogenic acid methyl ester) from this *R. carthamoides* along with 6 already reported constituents.

6 References

1. Biskup E, Lojkowska E. Evaluation of biological activities of *Rhaponticum carthamoides* extracts. JOURNAL OF MEDICINAL PLANTS RESEARCH. 2009 Dec;3(12):1092-8.
2. Kokoska L, Janovska D. Chemistry and pharmacology of *Rhaponticum carthamoides*: A review. Phytochemistry. 2009 May;70:842-55. <https://doi.org/10.1016/j.phytochem.2009.04.008>
3. Kolečkar V, Brojerova E, Rehakova Z, Kubikova K, Cervenka F, Kuca K, et al. In vitro antiplatelet activity of flavonoids from *Leuzea carthamoides*. Drug Chem Toxicol. 2008;31(1):27-35. <https://doi.org/10.1080/01480540701688444>
4. Maslov LN, Lishmanov YuB, Arbuzov AG, Krylatov A V, Budankova E V, Konkovskaya YuN, et al. Antiarrhythmic Activity of Phytoadaptogens in Short-Term Ischemia-Reperfusion of the Heart and Postinfarction Cardiosclerosis. Bull Exp Biol Med. 2009 Mar;147(3):331-4. <https://doi.org/10.1007/s10517-009-0502-6>
5. Peschel W, Kump A, Prieto JM. Effects of 20-hydroxyecdysone, *Leuzea carthamoides* extracts, dexamethasone and their combinations on the NF-kappa B activation in HeLa cells. JOURNAL OF PHARMACY AND PHARMACOLOGY. 2011 Nov;63(11):1483-95. <https://doi.org/10.1111/j.2042-7158.2011.01349.x>
6. Vyas AR, Singh S V. Molecular targets and mechanisms of cancer prevention and treatment by with-

- aferin a, a naturally occurring steroidal lactone. *AAPS J.* 2014;16(1):1-10. <https://doi.org/10.1208/s12248-013-9531-1>
7. Borovikova EB, Baitaev UA. Lesterone, a new phytoecdysteroid from the seeds of *Leuzea carthamoides*. *Chemistry of Natural Compounds.* 1999;35(2):205-6. <https://doi.org/10.1007/BF02234930>
 8. Sadykov ZT, Ramazanov NSh, Saatov Z. Phytoecdysteroids of Plants of The Genus *Rhaponticum*, Polypodin B 22-O-Benzooate from *Rhaponticum carthamoides*. 1997;33(6):665-6. <https://doi.org/10.1007/BF02249636>
 9. Ramazanov NSh, Makhimov ES, Saatov Z, Mamathkanov AU, Abdullaev ND. Phytoecdysteroids of plants of the genus *Rhaponticum* I. Carthamosterone a from Rh. *Carthamoides*. *Chemistry of Natural Compounds.* 1997;33(3):301-2. <https://doi.org/10.1007/BF02234880>
 10. Baltayev UA, Dinan L, Girault JP, Lafont R. 24(24(1)) [{}Z]-dehydroamarasterone B, a phytoecdysteroid from seeds of *Leuzea carthamoides*. *Phytochemistry.* 1997 Sep;46(1):103-5. [https://doi.org/10.1016/S0031-9422\(97\)00218-5](https://doi.org/10.1016/S0031-9422(97)00218-5)
 11. Buděšínský M, Vokac K, Harmatha J, Cvacka J. Additional minor ecdysteroid components of *Leuzea carthamoides*. *Steroids.* 2008 May;73:502-14. <https://doi.org/10.1016/j.steroids.2007.12.021>
 12. Girault JP, Lafont R, Varga E, Hajdu Z, Herke I, Szendrei K. Ecdysteroids from *Leuzea carthamoides*. *Phytochemistry.* 1988;27(3):737-41. [https://doi.org/10.1016/0031-9422\(88\)84085-8](https://doi.org/10.1016/0031-9422(88)84085-8)
 13. Glazowska J, Kaminski MM, Kaminski M. Chromatographic separation, determination and identification of ecdysteroids: Focus on Maral root (*Rhaponticum carthamoides*, *Leuzea carthamoides*). *J Sep Sci.* 2018 Dec;41(23):4304-14. <https://doi.org/10.1002/jssc.201800506>
 14. Pis J, Budesinsky M, Vokac K, Laudova V, Harmatha J. Ecdysteroids from the roots of *Leuzea carthamoides*. *Phytochemistry.* 1994 Oct;37(3):707-11. [https://doi.org/10.1016/S0031-9422\(00\)90343-1](https://doi.org/10.1016/S0031-9422(00)90343-1)
 15. Szendrei K, Varga E, Hajdu Z, Herke I, Lafont R, Girault JP. Ajugasterone-C and 5-deoxykaladasterone, an ecdysteroid artefact from *Leuzea carthamoides*. *J Nat Prod.* 1988;51(5):993-5. <https://doi.org/10.1021/np50059a034>
 16. Vokac K, Budesinsky M, Harmatha J. Minor ecdysteroid components of *Leuzea carthamoides*. *Collect Czechoslov Chem Commun.* 2002 Jan;67(1):124-39. <https://doi.org/10.1135/cccc20020124>
 17. Zhang XP, Zhang J, Dong M, Zhang ML, Huo CH, Shi QW, et al. Chemical Constituents of Plants from the Genus *Rhaponticum*. *Chem Biodivers.* 2010;7(3):594-609. <https://doi.org/10.1002/cbdv.200800275>
 18. Kokoska L, Janovska D. Chemistry and pharmacology of *Rhaponticum carthamoides*: A review. *Phytochemistry.* 2009 May 1;70(7):842-55. <https://doi.org/10.1016/j.phytochem.2009.04.008>
 19. Azizov AP, Seifulla RD. [The effect of elton, leveton, fitoton and adapton on the work capacity of experimental animals]. *Eksp Klin Farmakol.* 1998;61(3):61-63.
 20. Koudela K, Tenora J, Bajer J, Mathová A, Sláma K. Stimulation of growth and development in Japanese quails after oral administration of ecdysteroid-containing diet. *Eur J Entomol.* 1995;92:349-54.
 21. Kratky F, Hejhalek J, Kucharova S, Opletal L. Effect of 20-hydroxyecdysone on the protein synthesis in pigs. *Food and Agriculture Organisation of the United Nations.* 1998;42(10):445-51.
 22. Sláma K, Koudela K, Tenora J, Mathová A. Insect hormones in vertebrates: Anabolic effects of 20-hydroxyecdysone in Japanese quail. *Experientia.* 1996;52(7):702-6. <https://doi.org/10.1007/BF01925578>
 23. Syrov VN, Kurmukov AG. [Anabolic activity of phytoecdysone-ecdysterone isolated from *Rhaponticum carthamoides* (Willd.) Iljin]. *Farmakol Toksikol.* 1976;39(6):690-693.
 24. Gadzhieva RM, Portugalov SN, Paniushkin V V, Kondrat'eva II. [A comparative study of the anabolic action of ecdysten, leveton and Prime Plus, preparations of plant origin]. *Eksp Klin Farmakol.* 1995;58(5):46-48.
 25. Harmatha J, Vokác K, Kmoníčková E, Zídek Z. Lack of interference of common phytoecdysteroids with production of nitric oxide by immune-activated mammalian macrophages. *Steroids.* 2008 Apr;73(4):466-71. <https://doi.org/10.1016/j.steroids.2007.12.014>
 26. Lamer Zarawska E, Serafinowicz W, Gasirowski K, Brokos B. Immunomodulatory activity of polysaccharide-rich fraction from *Rhaponticum carthamoides* leaves. *Fitoterapia.* 1996;67(4):371-2.
 27. Logvinov S V, Pugachenko N V, Potapov A V, Krasnov EA, Plotnikov MB, Maslov MYu, et al. Ischemia-Induced Changes in Synptoarchitectonics of Brain Cortex and Their Correction with Ascovertin and *Leuzea* Extract. *Bull Exp Biol Med.* 2001;132(4):1017-20. <https://doi.org/10.1023/A:1013648118312>
 28. Petkov V, Roussinov K, Todorov S, Lazarova M, Yonkov D, Draganova S. Pharmacological Investigations on *Rhaponticum carthamoides*. *Planta Med.* 1984;50(03):205-9. <https://doi.org/10.1055/s-2007-969679>
 29. Mosharrof AH. Effects of extract from *Rhaponticum carthamoides* (Willd) Iljin (*Leuzea*) on learning and memory in rats. *Acta Physiol Pharmacol Bulg.* 1987;13(3):37-42.
 30. Plotnikov MB, Aliev OI, Vasil'ev AS, Maslov Mi, Dmitruk SE, Krasnov EA. Effect of *Rhaponticum carthamoides* extract on hemorheological properties of blood in rats with arterial hypertension. *Eksp Klin Farmakol.* 2001;64(6):45-47.
 31. Szendrei K, Reisch J, Varga E. Thiophene Acetylenes from *Leuzea* roots. *Phytochemistry.* 1984;23(4):901-2. [https://doi.org/10.1016/S0031-9422\(00\)85055-4](https://doi.org/10.1016/S0031-9422(00)85055-4)
 32. Huang Z, Dostal L, Rosazzas JPN. Mechanisms of Ferulic Acid Conversions to Vanillic Acid and Guaiacol by *Rhodotorula rubra*. *J Biol Chem.* 1993;268(32):23954-8. [https://doi.org/10.1016/S0021-9258\(20\)80477-3](https://doi.org/10.1016/S0021-9258(20)80477-3)
 33. Zhu X, Dong X, Wang Y, Ju P, Luo S. Phenolic Compounds from *Viburnum cylindricum*. *Helv Chim Acta [Internet].* 2005 Feb 1 [cited 2023 Jan 8];88(2):339-42. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1002/hlca.200590017> <https://doi.org/10.1002/hlca.200590017>
 34. Piš J, Buděšínský M, Vokáč K, Laudová V, Harmatha J. Ecdysteroids from the roots of *Leuzea carthamoides*. *Phytochemistry [Internet].* 1994 [cited 2023 Jan 8];37(3):707-11. Available from: https://www.academia.edu/10949508/Ecdysteroids_from_the_roots_of_Leuzea_carthamoides [https://doi.org/10.1016/S0031-9422\(00\)90343-1](https://doi.org/10.1016/S0031-9422(00)90343-1)

35. Buděšínský M, Vokáč K, Harmatha J, Cvačka J. Additional minor ecdysteroid components of *Leuzea carthamoides*. *Steroids*. 2008 May 1;73(5):502-14. <https://doi.org/10.1016/j.steroids.2007.12.021>
36. Ndongo JT, Mbing JN, Bikobo DN, Atchadé A de T, Shaaban M, Pegnyemb DE, et al. A new C-Glucosyl-flavone from *Sorindeia juglandifolia*. *Zeitschrift fur Naturforschung - Section C Journal of Biosciences* [Internet]. 2013 Jun 1 [cited 2023 Jan 8];68 C(5-6):169-74. Available from: <https://www.degruyter.com/document/doi/10.1515/znc-2013-5-601/html> <https://doi.org/10.1515/znc-2013-5-601>
37. Bastaev UA, Abubakirov NK. Phytoecdysteroids of *Rhaponticum carthamoides*. *Chem Nat Compd* [Internet]. 1987 [cited 2023 Jan 8];23:565-8. Available from: <https://agris.fao.org/agris-search/search.do?recordID=US8856798> <https://doi.org/10.1007/BF00598674>
38. Antonio Lemos da Silva L, Gabriela Faqueti L, Henrique Reginatto F, Diego Conceição dos Santos A, Barison A, Weber Biavatti M. Phytochemical analysis of *Vernonanthura tweediana* and a validated UPLC-PDA method for the quantification of eriodictyol. *Revista Brasileira de Farmacognosia* [Internet]. 2015 [cited 2023 Jan 8];25:375-81. Available from: <http://dx.doi.org/10.1016/j.bjp.2015.07.009>
39. Skiba A, Węglarz Z. Phenolic acids of *Rhaponticum carthamoides*. *Acta Hort.* 2004 Jan 30;597:119-24. <https://doi.org/10.17660/ActaHortic.2003.597.15>
40. Szendrei K, Varga E, Hajdú Z, Herke I, Lafont R, Girault JP. Ajugasterone C and 5-Deoxykaladasterone, an Ecdysteroid Artifact, from *Leuzea carthamoides*. *J Nat Prod*. 1988;51(5):993-5. <https://doi.org/10.1021/np50059a034>
41. Girault JP, Lafont R, Varga E, Hajdu Z, Herke I, Szendrei K. Ecdysteroids from *Leuzea carthamoides*. *Phytochemistry*. 1988;27(3):737-41. [https://doi.org/10.1016/0031-9422\(88\)84085-8](https://doi.org/10.1016/0031-9422(88)84085-8)
42. Wu TS, Yeh JH, Wu PL. The heartwood constituents of *Tetradium glabrifolium*. *Phytochemistry*. 1995 Sep 1;40(1):121-4. [https://doi.org/10.1016/0031-9422\(95\)00248-6](https://doi.org/10.1016/0031-9422(95)00248-6)
43. Dae SJ, Eun JP, Kang YH, Su BN, Hawthorne ME, Vigo JS, et al. Compounds obtained from *Sida acuta* with the potential to induce quinone reductase and to inhibit 7,12-dimethylbenz[*a*]anthracene-induced preneoplastic lesions in a mouse mammary organ culture model. *Arch Pharm Res* [Internet]. 2003 Aug 30 [cited 2023 Jan 8];26(8):585-90. Available from: <https://link.springer.com/article/10.1007/BF02976704> <https://doi.org/10.1007/BF02976704>
44. Chen JJ, Chung CY, Hwang TL, Chen JF. Amides and benzenoids from *Zanthoxylum ailanthoides* with inhibitory activity on superoxide generation and elastase release by neutrophils. *J Nat Prod*. 2009 Jan;72(1):107-11. <https://doi.org/10.1021/np800689b>
45. Park SJ, Kim N, Yoo G, Kim SN, Kwon HJ, Jung K, et al. Phenolics and neolignans isolated from the fruits of *Juglans mandshurica* Maxim. and their effects on lipolysis in adipocytes. *Phytochemistry*. 2017 May 1;137:87-93. <https://doi.org/10.1016/j.phytochem.2017.01.019>
46. Inoshiri S, Sasaki M, Kohda H, Otsum H, Yamasaki K. Aromatic glycosides from *Berchemia racemosa*. 1987;26(10):281-2814. [https://doi.org/10.1016/S0031-9422\(00\)83595-5](https://doi.org/10.1016/S0031-9422(00)83595-5)
47. Takara K, Otsuka K, Wada K, Iwasaki H, Yamashita M. 1,1-Diphenyl-2-picrylhydrazyl radical scavenging activity and tyrosinase inhibitory effects of constituents of sugarcane molasses. *Biosci Biotechnol Biochem*. 2007;71(1):183-91. <https://doi.org/10.1271/bbb.60432>
48. Luyen BTT, Tai BH, Thao NP, Yang SY, Cuong NM, Kwon YI, et al. A new phenylpropanoid and an alkylglycoside from *Piper retrofractum* leaves with their antioxidant and α -glucosidase inhibitory activity. *Bioorg Med Chem Lett*. 2014 Sep 1;24(17):4120-4. <https://doi.org/10.1016/j.bmcl.2014.07.057>
49. Wangenstein H, Miron A, Alamgir M, Rajia S, Samuelsen AB, Malterud KE. Antioxidant and 15-lipoxygenase inhibitory activity of rotenoids, iso-flavones and phenolic glycosides from *Sarcolobus globosus*. *Fitoterapia* [Internet]. 2006 [cited 2023 Jan 9];77(4):290-5. Available from: <https://pubmed.ncbi.nlm.nih.gov/16701962/> <https://doi.org/10.1016/j.fitote.2006.03.017>
50. Li T, Park MH, Kim MJ, Ryu BM, Kim MJ, Moon GS. Tachioside, an Antioxidative Phenolic Glycoside from Bamboo Species -Food Science and Biotechnology | Korea Science. *Food Sci Biotechnol* [Internet]. 2008 [cited 2023 Jan 8];17(6):1376-8. Available from: <https://koreascience.kr/article/JAKO200805441025627.page>